Brain Research, 495 (1989) 407–409
Elsevier

Author Index

Akiyama, H., see Tago, H., 271
Arnold, A.P., see Matsumoto, A., 362

Baran, H., Hörtnagl, H. and Hornykiewicz, O., Kainic acid-induced seizures: potentiation by α-methyl-*p*-tyrosine, 253
Baranyi, A., see Szente, M.B., 261
Barish, M.E., see Sands, S.B., 193
Barraclough, C.A., see Petersen, S.L., 9
Barth, T.M., see Stoessl, A.J., 20
Basinger, S.F., see Smiley, J.F., 31
Baty, L.T., see Gannon, R.L., 151
Beaulieu, C. and Colonnier, M., Effects of the richness of the environment on six different cortical areas of the cat cerebral cortex, 382
Besson, J.M., see Rivot, J.P., 140
Bickford-Wimer, P., see Boyajian, C.L., 66
Bidard, J.-N., Mourre, C., Gandolfo, G., Schweitz, H., Widmann, C., Gottesmann, C. and Lazdunski, M., Analogies and differences in the mode of action and properties of binding sites (localization and mutual interactions) of two K^+ channel toxins, MCD peptide and dendrotoxin I, 45
Bidard, J.-N., see Gandolfo, G., 189
Blake, P., see Kennaway, D.J., 349
Blaustein, J.D. and Turcotte, J.C., Estrogen receptor-immunostaining of neuronal cytoplasmic processes as well as cell nuclei in guinea pig brain, 75
Boyajian, C.L., Bickford-Wimer, P., Kim, M.B., Freedman, R. and Cooper, D.M.F., Pertussis toxin lesioning of the nucleus caudate-putamen attenuates adenylate cyclase inhibition and alters neuronal electrophysiological activity, 66
Broderick, P.A., Characterizing stearate probes in vitro for the electrochemical detection of dopamine and serotonin, 115
Bronstein, J., Wasterlain, C.G., Lasher, R. and Farber, D.B., Dark-induced changes in activity and compartmentalization of retinal calmodulin kinase in the rat, 83

Caramia, M.D., see Zarola, F., 217
Carpenter, D.O., see Evans, M.L., 309
Carpenter, D.O., see Salánki, J., 298
Chang, H.T. and Kuo, H., Calcitonin gene-related peptide (CGRP) in the rat substantia innominata and globus pallidus: a light and electron microscopic immunocytochemical study, 167
Chattarji, S., Stanton, P.K. and Sejnowski, T.J., Commissural synapses, but not mossy fiber synapses, in hippocampal field CA3 exhibit associative long-term potentiation and depression, 145
Colley, P.A., Sheu, F.-S. and Routtenberg, A., Dose-dependent phorbol ester facilitation or blockade of hippocampal long-term potentiation: relation to membrane/cytosol distribution of protein kinase C activity, 205
Colonnier, M., see Beaulieu, C., 382
Cooper, D.M.F., see Boyajian, C.L., 66
Crain, B., see Xie, C.-W., 156
Czéh, G. and Somjen, G.G., Spontaneous activity induced in isolated mouse spinal cord by high extracellular calcium and by low extracellular magnesium, 89

De Lanerolle, N.C., Kim, J.H., Robbins, R.J. and Spencer, D.D., Hippocampal interneuron loss and plasticity in human temporal lobe epilepsy, 387
Deutch, A.Y., see Knorr, A.M., 355
Douglass, J., see Xie, C.-W., 156
Dourish, C.T., see Stoessl, A.J., 20

Eastman, C.L. and Guilarte, T.R., Cytotoxicity of 3-hydroxykynurenine in a neuronal hybrid cell line, 225
Erulkar, S.D., see Stieber, A., 131
Evans, M.L. and Carpenter, D.O., Desensitization kinetics of a chloride acetylcholine response in *Aplysia*, 309
Evans, M.L., see Salánki, J., 298

Farber, D.B., see Bronstein, J., 83
Freedman, R., see Boyajian, C.L., 66
Frotscher, M., see Zipp, F., 161

Gandolfo, G., Gottesmann, C., Bidard, J.-N. and Lazdunski, M., Subtypes of K^+ channels differentiated by the effect of K^+ channel openers upon K^+ channel blocker-induced seizures, 189
Gandolfo, G., see Bidard, J.-N., 45
Ganon, R.L., Baty, L.T. and Terrian, D.M., L(+)-2-Amino-4-phosphonobutyrate inhibits the release of both glutamate and dynorphin from guinea pig but not rat hippocampal mossy fiber synaptosomes, 151
Gemba, H., see Sasaki, K., 100
Gonatas, N.K., see Stieber, A., 131
Gonias, S.L., see Hall, S.W., 373
Goping, G., see Nitsch, C., 243
Gottesmann, C., see Bidard, J.-N., 45
Gottesmann, C., see Gandolfo, G., 189
Greenwood, R.S., Meeker, R., Sullivan, H. and Hayward, J.N., Kindling in spontaneous hypertensive rats, 58
Grinnell, A.D., see Pawson, P.A., 182
Guilarte, T.R., see Eastman, C.L., 225

Hall, S.W., VandenBerg, S.R. and Gonias, S.L., Expression of plasminogen receptors on C6 glioma cells, 373
Hartman, R.D., see Petersen, S.L., 9
Hayward, J.N., see Greenwood, R.S., 58
Hersh, L.B., see Tago, H., 271
Hong, J.-S., see Xie, C.-W., 156
Hornykiewicz, O., see Baran, H., 253
Hörtnagl, H., see Baran, H., 253
Huang, L.-Y.M., Origin of thalamically projecting somatosensory relay neurons in the immature rat, 108
Hyde, T.M. and Peroutka, S.J., Distribution of cholecystokinin receptors in the dorsal vagal complex and other selected nuclei in the human medulla, 198

Iino, M., see Ozawa, S., 329
Ikeda, M., Mikuni, M., Nishikawa, T. and Takahashi, K., A neurochemical study of a new mutant mouse presenting myoclonus-like involuntary movement: a possible model of spontaneous serotonergic hyperactivity, 337

Isaacson, R.L., see Poplawsky, A., 396
Iversen, S.D., see Stoessl, A.J., 20

Jakobsen, L.J., see Raybould, H.E., 319
Jen, P.H.-S., see Sun, X., 1

Kato, T., see Xu, M., 232
Kennaway, D.J., Blake, P. and Webb, H.A., A melatonin agonist and *N*-acetyl-*N*²-formyl-5-methoxykynurenamine accelerate the reentrainment of the melatonin rhythm following a phase advance of the light–dark cycle, 349
Kim, J.H., see De Lanerolle, N.C., 387
Kim, M.B., see Boyajian, C.L., 66
Klatzo, I., see Nitsch, C., 243
Knorr, A.M., Deutch, A.Y. and Roth, R.H., The anxiogenic β-carboline FG-7142 increases in vivo and in vitro tyrosine hydroxylation in the prefrontal cortex, 355
Kudo, Y., see Ozawa, S., 329
Kuo, H., see Chang, H.T., 167

Lasher, R., see Bronstein, J., 83
Lazdunski, M., see Bidard, J.-N., 45
Lazdunski, M., see Gandolfo, G., 189
Leah, J. and Menetrey, D., Neuropeptides in propriospinal neurones in the rat, 173
Lee, P.H.K., see Xie, C.-W., 156

Marchand, R., see Woerly, S., 377
Mariorenzi, R., see Zarola, F., 217
Martin-Iverson, M.T., see Stoessl, A.J., 20
Martino, G., see Zarola, F., 217
Matsumoto, A., Arnold, A.P. and Micevych, P.E., Gap junctions between lateral spinal motoneurons in rats, 362
McGeer, E.G., see Tago, H., 271
McGeer, P.L., see Tago, H., 271
Meeker, R., see Greenwood, R.S., 58
Menetrey, D., see Leah, J., 173
Micevych, P.E., see Matsumoto, A., 362
Mikuni, M., see Ikeda, M., 337
Miller, L.P. and Perry, D.C., Opiate receptor subtype binding in gerbil hippocampus is altered by forebrain ischemia, 367
Miura, H., see Nakahara, D., 178
Miura, Y., see Nakahara, D., 178
Mizobe, F., see Xu, M., 232
Mourre, C., see Bidard, J.-N., 45

Nagatsu, T., see Nakahara, D., 178
Nakahara, D., Ozaki, Y., Miura, Y., Miura, H. and Nagatsu, T., Increased dopamine and serotonin metabolism in rat nucleus accumbens produced by intracranial self-stimulation of medial forebrain bundle as measured by in vivo microdialysis, 178
Nishikawa, T., see Ikeda, M., 337
Nitsch, C., Goping, G. and Klatzo, I., Preservation of GABAergic perikarya and boutons after transient ischemia in the gerbil hippocampal CA1 field, 243
Nitsch, R., see Zipp, F., 161
Novin, D., see Raybould, H.E., 319

Ogura, A., see Ozawa, S., 329
Ozaki, Y., see Nakahara, D., 178
Ozawa, S., Tsuzuki, K., Iino, M., Ogura, A. and Kudo, Y., Three types of voltage-dependent calcium current in cultured rat hippocampal neurons, 329

Paradiso, C., see Zarola, F., 217
Pawson, P.A. and Grinnell, A.D., Seasonal changes in the normal variability in release properties of motor nerve terminals in *Rana pipiens*, 182
Peroutka, S.J., see Hyde, T.M., 198
Perry, D.C., see Miller, L.P., 367
Petersen, S.L., Hartman, R.D. and Barraclough, C.A., An analysis of serotonin secretion in hypothalamic regions based on 5-hydroxytryptophan accumulation of push–pull perfusion. Effects of mesencephalic raphe or locus coeruleus stimulation and correlated changes in plasma luteinizing hormone, 9
Pointis, D., see Rivot, J.P., 140
Poplawsky, A. and Isaacson, R.L., Brief ganglioside treatment produces delayed enhancement of functional recovery after medial septal lesions, 396

Raybould, H.E., Jakobsen, L.J., Novin, D. and Taché, Y., TRH stimulation and L-glutamic acid inhibition of proximal gastric motor activity in the rat dorsal vagal complex, 319
Rivot, J.P., Pointis, D. and Besson, J.M., A comparison of the effects of morphine on 5-HT metabolism in the periaqueductal gray, ventromedial medulla and medullary dorsal horn: in vivo electrochemical studies in freely moving rats, 140
Robbins, R.J., see De Lanerolle, N.C., 387
Rossini, P.M., see Zarola, F., 217
Roth, R.H., see Knorr, A.M., 355
Routtenberg, A., see Colley, P.A., 205

Salánki, J., Evans, M.L. and Carpenter, D.O., Desensitization kinetics of a K^+ acetylcholine response in *Aplysia*, 298
Sands, S.B. and Barish, M.E., NMDA receptor activation by residual glutamate in glutamine preparations: a cautionary note regarding weak NMDA receptor agonists, 193
Sasaki, K., Gemba, H. and Tsujimoto, T., Suppression of visually initiated hand movement by stimulation of the prefrontal cortex in the monkey, 100
Schweitz, H., see Bidard, J.-N., 45
Sejnowski, T.J., see Chattarji, S., 145
Senba, E., see Tamatani, M., 122
Sheu, F.-S., see Colley, P.A., 205
Smiley, J.F. and Basinger, S.F., Glycine high-affinity uptake labels a subpopulation of somatostatin-like immunoreactive cells in the *Rana pipiens* retina, 31
Somjen, G.G., see Czéh, G., 89
Soriano, E., see Zipp, F., 161
Spencer, D.D., see De Lanerolle, N.C., 387
Stanton, P.K., see Chattarji, S., 145
Stewart, J. and Vezina, P., Microinjections of Sch-23390 into the ventral tegmental area and substantia nigra pars reticulata attenuate the development of sensitization to the locomotor activating effects of systemic amphetamine, 401
Stieber, A., Erulkar, S.D. and Gonatas, N.K., A hypothesis for the superior sensitivity of wheat germ agglutinin as a neuroanatomical probe, 131
Stoessl, A.J., Martin-Iverson, M.T., Barth, T.M., Dourish, C.T. and Iversen, S.D., Effects of ageing on the behavioural responses to dopamine agonists: decreased yawning and locomotion, but increased stereotypy, 20
Sullivan, H., see Greenwood, R.S., 58
Sun, D., see Sun, X., 1
Sun, X., Jen, P.H.-S., Sun, D. and Zhang, S., Corticofugal influences on the responses of bat inferior colliculus neurons to sound stimulation, 1

Szente, M.B. and Baranyi, A., Properties of depolarizing plateau potentials in aminopyridine-induced ictal seizure foci of cat motor cortex, 261

Taché, Y., see Raybould, H.E., 319
Tago, H., McGeer, P.L., McGeer, E.G., Akiyama, H. and Hersh, L.B., Distribution of choline acetyltransferase immunopositive structures in the rat brainstem, 271
Takahashi, K., see Ikeda, M., 337
Tamatani, M., Senba, E. and Tohyama, M., Calcitonin gene-related peptide- and substance P-containing primary afferent fibers in the dorsal column of the rat, 122
Terrian, D.M., see Gannon, R.L., 151
Tohyama, M., see Tamatani, M., 122
Traversa, R., see Zarola, F., 217
Tsujimoto, T., see Sasaki, K., 100
Tsuzuki, K., see Ozawa, S., 329
Turcotte, J.C., see Blaustein, J.D., 75

VandenBerg, S.R., see Hall, S.W., 373
Vezina, P., see Stewart, J., 401

Wasterlain, C.G., see Bronstein, J., 83
Webb, H.A., see Kennaway, D.J., 349
Widmann, C., see Bidard, J.-N., 45
Woerly, S. and Marchand, R., A fluorescence histochemical method for the demonstration of central catecholamine neurons in young embryonic tissues, 377

Xie, C.-W., Lee, P.H.K., Douglass, J., Crain, B. and Hong, J.-S., Deep prepyriform cortex kindling differentially alters the level of prodynorphin mRNA in rat hippocampus and striatum, 156
Xu, M., Mizobe, F., Yamamoto, T. and Kato, T., Differential effects of M_1- and M_2-muscarinic drugs on striatal dopamine release and metabolism in freely moving rats, 232

Yamamoto, T., see Xu, M., 232

Zarola, F., Caramia, M.D., Paradiso, C., Mariorenzi, R., Martino, G., Traversa, R. and Rossini, P.M., Single fibre motor evoked potentials to brain, spinal roots and nerve stimulation. Comparisons of the 'central' and 'peripheral' response jitter to magnetic and electric stimuli, 217
Zhang, S., see Sun, X., 1
Zipp, F., Nitsch, R., Soriano, E. and Frotscher, M., Entorhinal fibers form synaptic contacts on parvalbumin-immunoreactive neurons in the rat fascia dentata, 161